



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Leontodon nudicaule of Jelliffe's list is evidently based on a specimen of Dr. Hulst's so labeled which specimen is

Picris hieracioides L.

Lactuca sagittifolia Ell. Rockaway.

Crepis virens L. Richmond Hill, Cypress Hills, *Hulst.*

Xanthium Canadense Mill. The common form is var. *echinatum*.

Solidago juncea Ait. Cypress Hills, *Hulst.*

Hypochaeris radicata L. Cedarhurst.

Antennaria neglecta Greene. Common in sandy fields.

Senecio obovatus Muhl. Richmond Hill, *Hulst.*

BOYS' HIGH SCHOOL, BROOKLYN.

A KEY TO THE NORTH AMERICAN GENERA AND SPECIES OF THE HYGROPHOREAE.—I

BY F. S. EARLE

KEY TO THE GENERA

- | | |
|---|------------------------|
| 1. Spores black or dark brown. | A. GOMPHIDIUS. |
| Spores white or whitish. | 2. |
| 2. Usually parasitic on other agarics; basidiospores mostly replaced by chlamydospores. | B. NYCTALIS. |
| Not parasitic on agarics; no chlamydospores. | 3. |
| 3. With a glutinous veil when young, leaving a more or less persistent annulus. | C. LIMACIUM. |
| With no veil or annulus. | D. HYGROPHORUS. |

A. KEY TO THE NORTH AMERICAN SPECIES OF GOMPHIDIUS

- | | |
|---|---------------------------------|
| 1. Pileus white or whitish. | 2. |
| Pileus some shade of red or brown. | 3. |
| Pileus dingy pink; lamellae whitish; stipe yellow. | <i>G. flavipes</i> Pk. |
| 2. Stipe yellow, shorter than the diameter of the pileus. | <i>G. maculatus</i> (Scop.) Fr. |
| Stipe white, longer than the diameter of the pileus. | <i>G. furcatus</i> Pk. |
| 3. Stipe concolorous (reddish or brownish). | 4. |
| Stipe at first white or whitish. | 7. |

4. Entire plant blackening in drying. *G. Oregonensis* Pk.
Plant not blackening, or only the gluten blackening. 5.
5. Plant not at all blackening ; stipe more or less tomentose. 6
The gluten on pileus blackening ; stipe pruinose at base. *G. vinicolor* Pk.
6. Large, 5-8 cm. ; stipe longer than diameter of pileus, thinly brown-tomentose.
G. viscidus (L.) Fr.
Smaller, 2-4 cm. ; stipe shorter than diameter of pileus, yellow-tomentose below,
glabrate above. *G. Alabamensis* Earle.
7. Entire plant turning jet black in drying. *G. nigricans* Pk.
Plant not blackening. 8.
8. Pileus purplish brown ; stipe white, yellow within. *G. glutinosus* (Schaeff.) Fr.
Pileus rose-color ; stipe white, base and interior flesh-color. *G. roseus* Fr.

B. KEY TO THE NORTH AMERICAN SPECIES OF NYCTALIS

1. Parasitic on *Russula*, etc. *N. asterophora* Fr.

C. KEY TO THE NORTH AMERICAN SPECIES OF LIMACIUM

1. Pileus white or whitish, disc often colored. 2.
Pileus grayish yellow. *L. elegantulum* (Pk.)
Pileus with some shades of blue or violet. 7.
Pileus grayish brown or blackish brown. 8.
2. Pileus with more or less yellow on disc. 4.
Pileus with brownish disc and innate black fibrils. *L. virgatum* (Pk.) P. Henn.
Pileus white throughout. 3.
3. Spores irregular-sphaeroid, 5-6 μ . *L. eburneum* (Bull.) Schroet.
Spores ellipsoid, 6.5-7.5 $\mu \times$ 4-5 μ . *L. sordidum* (Pk.)
4. Lamellae stained greenish yellow with age ; pileus covered with yellow gluten.
L. paludosum (Pk.)
Lamellae becoming flesh-color ; disc yellow or reddish. *L. flavo-discum* (Frost)
Lamellae white, unchanging. 5.
5. Pileus with deciduous yellow flecks on the disc and lamellae.
L. chrysodon (Batsch) Schroet.
Pileus smooth, glutinous. 6.
6. Stipe white, farinose above. *L. luteum* (Johns.)
Stipe yellowish white, apex roughened. *L. Laurae* (Morg.) P. Henn.
7. Pileus bluish-alutaceous, lamellae darker. *L. caerulescens* (B. & C.)
Pileus violaceous when moist, paler when dry, lamellae violaceous.
L. subviolaceum (Pk.)
8. Margin of lamellae entire ; stipe white or brown-spotted.
L. fuligineum (Frost) P. Henn.
Margin of lamellae erose ; stipe pallid or brownish. *L. Morrisii* (Pk.)